

Bell's Inequality and τ -Physics at LEP

H. Dreiner

Oct. 29th, 1992

Theoretical Physics, University of Oxford,
1 Keble Road, Oxford OX1 3NP, England

Abstract

Abstract

In this talk presented at the TAU92 workshop, Columbus, OH, Sept. 92, we summarize results presented in more detail in a recent paper by S. Abel, M. Dittmar and the author where we gave a general proof that Bell's inequality can not be tested at a collider experiment. In particular, a measurement of correlated tau-spins at LEP does not constitute a test of local realistic theories via Bell's inequality. The central point of the argument is that such tests, where the spins of two particles are inferred from a scattering distribution, can always be described by a local hidden variable theory. In response to questions at the workshop we go beyond the paper and show that an old experiment involving the measurement of the correlated spins of the two photons emitted in positronium decay via Compton-scattering is also not a viable test of Bell's inequality.

1 Introduction:

Since the 1935 paper [1] of Einstein, Podolsky, and Rosen (EPR) the question of whether quantum mechanics (QM) offers a complete description of “physical reality” has received much attention. Since the work of Bell [2, 3], this question has been under the scrutiny of experiment. Bell's theorem states that in all local-realistic theories [4] two-particle correlation functions satisfy Bell's inequality, whereas the corresponding quantum mechanical correlation functions can violate the inequality in some regions of parameter space. To date there are many experimental tests of Bell's inequality [5]-[11] which we find worth complementing in three respects [13]:

- We think it would be of interest to test the inequalities with fermions and/or with massive particles; fermions are peculiar to QM and a test with massive particles could exclude certain hidden-variable theories.
- The correlation in the atomic cascade experiment is physically established through electromagnetic interactions; there have been profound surprises in the physics of

the weak-interactions, notably P- and CP-violation, and perhaps we are in store for another.

- The energy scale probed in the atomic cascade experiments is in the eV range; as Bohm has pointed out, non-local effects may well become apparent at a length-scale below about $10^{-13} cm$ [12]. It is thus important to test for non-local features at as high an energy-scale as possible.

In a recent paper [13] we showed that it is *not* possible to test locality via Bell's inequality in a collider experiment. It is therefore not possible to devise a novel test involving the weak interactions at LEP for example. We summarize the argument here.

2 Proposed Experimental Test at LEP

As a possible test of Bell's inequality with massive fermions we consider τ -production at LEP, $e^+e^- \rightarrow Z^0 \rightarrow \tau^+\tau^-$. The two taus in the Z^0 -decay are produced with opposite chirality and thus with opposite helicity ($M_{Z^0} \gg m_\tau$). Since the τ 's have equal and opposite momentum in the lab-frame, the τ -spins along their momentum axes are *positively* correlated.

This system of two spin- $\frac{1}{2}$ particles with positively correlated spins is to be considered for an experimental test of Bell's inequality. It has been shown at LEP that the helicity of the tau can be determined with good accuracy from the momentum spectrum of the decay products [14]. To simplify the discussion we concentrate on the decay $\tau \rightarrow \pi\nu_\tau$.

The cross-section for the process $e^+e^- \rightarrow \pi^+\pi^-\bar{\nu}_\tau\nu_\tau$ calculated in the Standard Model is

$$\frac{d\sigma}{d\cos\theta_{\pi\pi}}(e^+e^- \rightarrow \pi^+\pi^-\bar{\nu}_\tau\nu_\tau) = A(1 - \frac{1}{3}\cos\theta_{\pi\pi}), \quad (1)$$

where the factor A is given in [13] and $\cos\theta_{\pi\pi} = (\hat{p}_{\pi^+} \cdot \hat{p}_{\pi^-})$. The pion momenta ($\hat{p}_{\pi^\pm}$) are given in the respective parent τ rest-frames.

3 Testing Bell's Inequality?

In the Standard Model the expectation value for observing two pions at relative angle $\cos\theta_{\pi\pi}$ is given by

$$P_{QM}(\cos\theta_{\pi\pi}) = \frac{\frac{d\sigma}{d\cos\theta_{\pi\pi}}(e^+e^- \rightarrow \pi^+\pi^-\bar{\nu}_\tau\nu_\tau)}{\sigma(e^+e^- \rightarrow \pi^+\pi^-\bar{\nu}_\tau\nu_\tau)} \quad (2)$$

$$= \frac{1}{2}(1 - \frac{1}{3}\cos\theta_{\pi\pi}). \quad (3)$$

A more general form of Bell's inequality than given in [13], which also takes into account detector efficiencies is given in [15]

$$2 - |P(\cos\theta_{\hat{a}\hat{c}}) + P(\cos\theta_{\hat{a}\hat{d}})| \geq |P(\cos\theta_{\hat{b}\hat{c}}) - P(\cos\theta_{\hat{b}\hat{d}})| \quad (4)$$

When inserting the form (3) we obtain

$$12 - |6 - \cos \theta_{\hat{a}\hat{c}} - \cos \theta_{\hat{a}\hat{d}}| \geq |\cos \theta_{\hat{b}\hat{c}} - \cos \theta_{\hat{b}\hat{d}}|, \quad (5)$$

which is satisfied for all sets angles $\theta_{\hat{x}\hat{y}}$. Here $\theta_{\hat{x}\hat{y}}$ denotes the angle between two pions emitted in directions $\hat{x}$ and $\hat{y}$ respectively. We thus conclude that as presented this experiment is not a test of Bell's inequality.

It is worth pointing out that this conclusion does not directly depend on the factor $\frac{1}{3}$ in Eq.(3). If we consider a more general form $P(\cos \theta) = \frac{1}{2}(1 - a \cos \theta)$, we find after insertion into Eq.(4) that it is satisfied for all $a \leq 1$:

$$\frac{4}{a} - \left| \frac{2}{a} - a \cos \theta_{\hat{a}\hat{c}} - a \cos \theta_{\hat{a}\hat{d}} \right| \geq |\cos \theta_{\hat{b}\hat{c}} - \cos \theta_{\hat{b}\hat{d}}|. \quad (6)$$

4 Conceptual Deficiency of the Experiment

It is straightforward, using a construction proposed by Kasday in a different context [6], to construct an *ad hoc* local hidden variable theory LHVT which also leads to the result (3) for the production of charged pions via τ -pairs at LEP. The construction goes as follows: the Standard Model (and thus QM) provides the differential cross-section which is a function of the unit pion momenta¹

$$\frac{d\sigma}{d\Omega_+ d\Omega_-}(e^+e^- \rightarrow \pi^+\pi^-\nu_\tau\bar{\nu}_\tau) = f(\hat{p}_{\pi^+}, \hat{p}_{\pi^-}), \quad (7)$$

where as before we consider the pion momenta in the respective rest frames of the tau-leptons; and $d\Omega_\pm$ denotes the angular phase space of the two pions respectively. Now let the hidden variables for each tau-lepton be a set of unit vectors $(\hat{\lambda}_e, \hat{\lambda}_\mu, \hat{\lambda}_\pi, \hat{\lambda}_\rho, \dots)$. When the tau decays as $\tau^- \rightarrow \pi^-\nu_\tau$, $\hat{\lambda}_\pi$ tells it to decay such that the pion momentum is in the direction $\hat{p}_\pi = \hat{\lambda}_\pi$. Now let $F(\hat{\lambda}_{\pi^+}, \hat{\lambda}_{\pi^-})$ be the original probability distribution of the hidden variables. If we identify this LHVT with the QM function of Eq.(7)

$$F(\hat{\lambda}_{\pi^+}, \hat{\lambda}_{\pi^-}) = f(\hat{\lambda}_{\pi^+}, \hat{\lambda}_{\pi^-}), \quad (8)$$

we obtain a hidden-variable theory which will exactly reproduce all experimental results of the process $e^+e^- \rightarrow \pi^+\pi^-\nu_\tau\bar{\nu}_\tau$.

It is essential in the above construction that QM provides the function $f(\hat{p}_{\pi^+}, \hat{p}_{\pi^-})$. This in turn is only possible because all the components of $\hat{p}_{\pi^\pm}$ commute with each other

$$\begin{aligned} [(\hat{p}_{\pi^+})_i, (\hat{p}_{\pi^-})_j] &= 0, \\ [(\hat{p}_{\pi^\pm})_i, (\hat{p}_{\pi^\pm})_j] &= 0. \end{aligned} \quad (9)$$

By contrast, when measuring non-commuting operators in the final state this construction is no longer possible. For example, in the two photon experiments [7], when measuring

¹ From $f(\hat{p}_{\pi^+}, \hat{p}_{\pi^-})$ we obtained Eq.(1) by integrating over all but the relative angle between the pions. The formula for $f(\hat{p}_{\pi^+}, \hat{p}_{\pi^-})$ is very extensive and we do not include it here. It should however be clear that such a function exists and this is all that we require.

the transverse spin components S_a^i , $i = 1, 2$, $a = x, y$, of the two correlated photons we have

$$[S_a^1, S_b^2] = 0, \quad [S_a^i, S_b^i] = i\hbar\epsilon_{abz}S_z^i. \quad (10)$$

In this case QM does *not* provide a function $f(S_x^i, S_y^i, \dots)$ because QM can not give a function for an experimental result which depends on the simultaneous knowledge (measurement) of two non-commuting variables. Therefore the Kasday construction breaks down.

We conclude that in all experiments where the correlated observables commute, it is possible using the method of Eq.(8) to construct a LHVT which reproduces the QM results and thus the experimental results. In these cases the QM result must satisfy Bell's inequality just like the LHVT result does. In the case of an alternative experiment which measures non-commuting observables this is not the case and we are left with the *possibility* that it is a viable test of Bell's inequality. Thus non-commuting observables are a *necessary* requirement for a valid test of locality.

Our conclusion similarly applies to the proton fixed target experiment [9]; the Λ pair production experiment [10, 11] and the positronium decay experiments [8], which we return to below.

5 Testing Hidden Variable Theories

Despite the above criticism several experiments involving commuting observables have been performed [8]-[11]. We briefly outline the method employed and the assumptions made in order to extract a correlation function from the experiments which does violate Bell's inequality. We go on to outline why we disagree with this method.

Given the correlation $P_{\pi\pi}(\theta) = \frac{1}{2}(1 - \frac{1}{3}\cos\theta_{\pi\pi})$, it is clear if we consider the function

$$P(\theta) = 3(P_{\pi\pi}(\theta) - \frac{1}{2}) = -\cos\theta \quad (11)$$

we find that for special sets of angles $P(\theta)$ violates Bell's inequality (4). Presented in this form $P(\theta)$ has no physical interpretation and it is thus irrelevant whether it violates Bell's inequality or not. Recall that the objective of experimental tests of Bell's inequality is to test whether nature is described by a local theory or by a non-local theory such as QM. We now proceed to show that *if* one assumes that QM is the proper description (despite the fact that we want to test it!) *then* $P(\theta)$ can be given a physical interpretation and given this strong assumption one can "test Bell's inequality".

Within the framework of QM one can derive [13] on symmetry grounds a general formula for the expectation value of the relative angle between the final state pions emitted from the spin- $\frac{1}{2}$ τ 's

$$P_{\pi\pi}(\theta) = c_1 + c_2 < (\hat{p}_{\pi^+} \cdot \vec{\sigma}_{\tau^+})(\hat{p}_{\pi^-} \cdot \vec{\sigma}_{\tau^-}) >_{QM}. \quad (12)$$

for general constants c_1 and c_2 . This corresponds to s - and p -wave scattering. The second term on the right-hand side is the expectation value for the τ^+ -spin to be observed in

the direction $\hat{p}_{\pi^+}$ and the τ^- -spin in the direction $\hat{p}_{\pi^-}$. But keep in mind that this is all within the framework of QM. This second term has an exact analogon in the better known atomic cascade experiments [7], where the transverse spins of two photons, γ_a, γ_b , in the singlet state are measured by standard polarizers set-up in (unit-vector) directions $\hat{a}$ and $\hat{b}$. The expectation value that the two photons are observed is then given by

$$P_{\gamma_a \gamma_b} = \langle (\hat{a} \cdot \vec{\sigma}_{\gamma_a})(\hat{b} \cdot \vec{\sigma}_{\gamma_b}) \rangle_{QM}, \quad (13)$$

$$= -\hat{a} \cdot \hat{b}. \quad (14)$$

This expectation value violates Bell's inequality for certain sets of angles. We can see from Eq.(12) that within the framework of QM the expectation value of the correlation of the τ -spins

$$P_{\sigma_\tau \sigma_\tau}(\theta) = \langle (\hat{a} \cdot \vec{\sigma}_{\tau^+})(\hat{b} \cdot \vec{\sigma}_{\tau^-}) \rangle_{QM} \quad (15)$$

can be given in terms of the expectation value of the correlated pion momenta

$$P_{\sigma_\tau \sigma_\tau}(\theta) = \frac{P_{\pi\pi}(\theta) - c_1}{c_2} = -\cos(\theta), \quad (16)$$

which is just $P(\theta)$ of Eq.(11). If we now make the in our opinion strong assumption that this is true in general, *independently* of whether QM describes the correlation correctly or not (remember that is what we are trying to check) and eventhough we used QM to obtain Eq.(16), then we can experimentally test $P_{\sigma_\tau \sigma_\tau}(\theta)$ and the correlation of the tau spins by measuring $P_{\pi\pi}(\theta)$. As given above, $P_{\sigma_\tau \sigma_\tau}(\theta)$ violates Bell's inequality and is thus not reproduceable by any LHVT.

It is tempting to conclude that this is a test of locality via Bell's inequality. We argue instead that it is merely a test of a subclass of LHVTs, where the size of the subclass is unknown. In obtaining Eq.(16) we have used QM to translate our expectation of the pions ($P_{\pi\pi}$) into an expectation about the tau-spins ($P_{\tau\tau}$). Thus we are assuming QM to start with. Dynamically the connection is given through the tau-decay and we are therefore specifically assuming QM (*i.e.* the SM) describes the tau-decay properly *as well as* the correlations we establish or maintain through it. However, our main objective is to test whether QM offers a complete description of nature, and thus it is inconsistent to employ it in the test. If we do make the assumption Eq.(16) or Eq.(12) then of course we are restricting ourselves to a test of only those LHVTs which also satisfy Eq.(12). Thus we are testing a subclass of LHVTs, but there is a subclass of LHVTs of unknown size which remains untested. The LHVT we constructed in section 4 for example falls into the second category.

One might now reply that we know from measurements on single taus that QM describes the tau decay correctly. Indeed we do, but it is also known that it is trivial to construct a LHVT which correctly describes the decay of a single tau as well. The distinction between naïve local-realistic expectations (LHVTs) and QM doesn't arise until we consider an experiment involving the correlation of at least two particles. This point is at the heart of the EPR paradox as well as the work of Bell. Thus it is exactly at the level of the established correlation that we want to test QM *versus* LHVTs and it

is not appropriate to assume the correctness of QM in order to consider an *unobserved* correlation.

The above argument might be considered to be implicit in Kasday's paper [6]. However, in the subsequent literature it has been ignored [9, 10, 11, 16, 17]. We have thus found it important to stress and elucidate the full extent of the argument.

6 Compton Scattering Experiment

In response to discussions at the workshop we conclude with a brief review of an old experimental test of Bell's inequality where polarization correlations are observed for pairs of photons emitted in the decay of positronium [8]. From parity considerations it is known that the photons are in the singlet state. The spins of the 500 keV photons are "analyzed" via Compton scattering, and the correlation is thus an angular correlation in the momenta of the scattered photons. This is thus an experiment where commuting observables are measured and from our proof we conclude that the experiment is *not a test of Bell's inequality*.

This can also be seen explicitly from the QM formula for the scattering correlation. The correlation is given by the Klein-Nishina Compton scattering formula as for example given in [4]

$$P(\theta_1, \phi_1, \theta_2, \phi_2) = \frac{1}{16\pi^2} (1 - C \frac{G(\theta_1)G(\theta_2)}{F(\theta_1)F(\theta_2)} \cos 2(\phi_1 - \phi_2)) \quad (17)$$

where we have introduced a normalization factor as in Eq.(2) and

$$F(\theta) = [2 + (1 - \cos \theta)^3] / (2 - \cos \theta)^3, \quad (18)$$

$$G(\theta) = \sin^2 \theta / (2 - \cos \theta)^2. \quad (19)$$

The constant $C \leq \frac{1}{2}$ and θ_i, ϕ_i are the final state angles of the photons $i = 1, 2$. As an example consider $P(\theta_1, \phi_1, \theta_2, \phi_2)$ as a function only of the relative angle $\Delta\phi = \phi_1 - \phi_2$ and with fixed θ_1, θ_2

$$P(\Delta\phi) = \frac{1}{\pi} (1 - C \frac{G(\theta_1)G(\theta_2)}{F(\theta_1)F(\theta_2)} \cos 2(\Delta\phi)), \quad (20)$$

It is straightforward to show that $G(\theta)/F(\theta) < 1$ for all θ and thus using the argument of Eq.(6) Section 3 we see that $P(\Delta\phi)$ satisfies Bell's inequality for any angles $\Delta\phi, \theta_1, \theta_2$.

We also point out, that Bell constructed a LHVT [18] which gives *exactly* the same prediction for the correlation of the two scattered photons as the QM prediction given in Eq.(17). The statement of Bell's theorem [2] is that *every* LHVT must satisfy Bell's inequality. Thus the LHVT constructed by Bell and the identical QM prediction given in Eq.(17) must satisfy Bell's inequality for *all* angles $\theta_1, \phi_1, \theta_2, \phi_2$. And we conclude that this experiment is not a viable test of Bell's inequality.

Acknowledgements The author would like to thank Steven Abel and Michael Dittmar with whom this work was done.

References

- [1] A. Einstein, B. Podolski and N. Rosen, Phys. Rev. 37 (1935) 777.
- [2] J.S. Bell, Physics 1 (1964) 195.
- [3] J.S. Bell, Rev. Mod. Phys. 38 (1966) 447.
- [4] For a review of hidden-variable theories see: F.J. Belinfante, *A Survey of Hidden-Variable Theories*, Pergamon Press (1973).
- [5] C.A. Kocher and E.D. Commins, Phys. Rev. Lett. 18 (1967) 575; C. Papaliolios, Phys. Rev. Lett. 18 (1967) 622; C. Dewdney *et al*, Phys. Lett. 113A (1985) 135; D.R. Inglis, Rev. Mod. Phys. 33 (1961) 1; J. Six, Phys. Lett. 114B (1982) 200; F. Selleri, Lett. Nuovo Cim. (1983) 521; A. Datta, D. Home and A. Raychaudhuri, Phys. Lett. 123A (1987) 4; T.K. Lo and A. Shimony, Phys. Rev. A23 (1981) 3003; E.Santos, Phys. Rev. A30 (1984) 2128.
- [6] L.R.Kasday, in Proc. Int. School of Physics ‘Enrico Fermi’, course 49 (1971) p.195.
- [7] S.J. Freedman and J.F. Clauser, Phys. Rev. Lett. 28 (1972) 938; E.S. Fry, and R.C. Thompson, Phys. Rev. Lett. 37 (1976) 465; A. Aspect and P. Grangier, in Proc. Int. Symp. Foundations of Quantum Mechanics, Tokyo (1983) 214; A. Aspect, and P. Grangier, Lett. Nuovo Cim. 43 (1985) 345; A. Aspect, J. Dalibard and G. Roger, Phys. Rev. Lett. 49 (1982) 1804; J.F. Clauser, M.A. Horne, A. Shimony and R.A. Holt, Phys. Rev. Lett. 23 (1969) 880.
- [8] E. Bleuler and H.L. Brandt, Phys. Rev. 73 (1948) 1398; C.S. Wu and J. Shalnov, Phys. Rev. 77 (1950) 136; H. Langhoff, Z. Phys. 160 (1960) 186; L.R. Kasday, J. Ullman and C.S. Wu, Bulletin Am. Phys. Soc. 15 (1970) 586.
- [9] M. Lamehi-Rachti and W. Mittig, Phys. Rev. D14 (1976) 2543.
- [10] N.A. Törnqvist, Found. Phys. 11 (1981) 171; Europhys. Lett. 1 (1986) 381; Phys. Lett. 117A (1986) 1.
- [11] M.H.Tixier *et al*, DM2 collaboration, Phys. Lett. B 212(1988) 523.
- [12] D. Bohm, Phys.Rev. 85(1952)166, 180.
- [13] S. Abel, M. Dittmar, and H. Dreiner, Phys. Lett. B 280 (1992) 304.
- [14] Aleph collaboration, D. Decamp *et al*, Phys. Lett. B265 (1991) 430; Opal collaboration, Phys. Lett. B266 (1991) 201.
- [15] Eq.(226), Part III, Section 4.9 of Ref.[4].
- [16] *Quantum Theory and Measurement*, edited by J.A. Wheeler and W.H. Zurek, Princeton Series and Physics, Princeton Univ. Press, 1983.

- [17] P. Privitera, Phys. Lett. B 275 (1992)172.
- [18] See Eqs.(G-1a) and (G-1b) of Appendix G in Ref.[4].